



COVID-19 Management Protocol AIIMS, New Delhi

20th March 2020

COVID-19 Suspect

- i) Any patient with acute respiratory illness (fever with at least one of the following- cough or shortness of breath) with:
- History of travel to high-risk COVID-19 affected countries in the last 14 days, or
 - Close contact with a laboratory confirmed case of COVID-19 in the 14 days, or
 - Health care personnel (HCP) managing respiratory distress/severe acute respiratory illness cases, when they are symptomatic

- Asymptomatic traveller/close contact**
- Home quarantine
 - Twice daily self monitored temperature
 - Contact & droplet precautions

On developing symptoms

Mild case

- Low-grade fever, cough, malaise, rhinorrhea, sore throat without shortness of breath
- Treatment**
 Tab oseltamivir 75mg BD (for high-risk influenza suspects)
 Antibiotics if needed (azithromycin+ amox /clav)
 Tab Paracetamol 500 mg SOS
 Symptomatic

Moderate to severe case

- Any one of:**
1. Respiratory rate > 24/min
 2. SpO₂ < 94% in room air
 3. Confusion/drowsiness
 4. Systolic BP < 90 mmHg or diastolic BP < 60 mmHg

Admit & test

- Test negative**
- Manage according to existing protocol

- Test positive**
- Oxygen supplementation to maintain SpO₂>94%
 - Antipyretics, antitussives, antibiotics as indicated
 - MDI preferred over nebulization
 - Hydroxychloroquine (400 mg BD x 1 day f/b 200 mg BD x 5 days) may be considered
 - Lopinavir/ritonavir(200 mg 2 tab BD) may be considered on case-to-case basis (within 10 days of symptom-onset)
 - Do not combine Hydroxychloroquine with Lopinavir in view of drug interactions
 - Corticosteroids to be avoided

- Test negative**
- Symptomatic management

- Test positive**
- Home isolation (>72 hrs afebrile or 7 days after symptom onset whichever is longer)/two negative samples 24 hours apart
 - Self-monitoring for fever
 - Paracetamol & symptomatic Rx
 - Contact & droplet precautions
 - Danger signs explained
 - High-risk individuals* may be considered for admission based on clinical judgement

Call helpline
011- 23978046

- If worsening**
- Respiratory failure
 - Hypotension
 - Worsening mental status
 - MODS

Shift to ICU

- NIV/HFNC to be used carefully in view of risk of aerosol generation
- Ventilator management as per ARDS protocol
- Conservative fluid management (if not in shock)
- Standard care for ventilated patient
- Closed suction and HME filters
- Prone ventilation, ECMO for refractory hypoxemia.

After clinical & radiological improvement

Discharge
if two negative samples at least 24 hours apart

Improving

- *High-risk for severe disease**
- ✓ Age > 60 years
 - ✓ Cardiovascular disease including hypertension
 - ✓ DM, other immunocompromised states
 - ✓ Chronic lung/kidney/liver disease